



Khasi Hills Community REDD+ Project: Restoring and Conserving Meghalaya's Hill Forests through Community Action



REDD+ and ANR Technical Specifications

Submitted to Plan Vivo, UK by

Community Forestry International on behalf of

Ka Synjuk Ki Hima Arliang Wah Umiam,

Mawphlang Welfare Society

Mawphlang, Meghalaya, North Eastern India



VERSION 4.0
15 November,
2018



SUMMARY OF PROJECT

PROJECT TITLE	Khasi Hills REDD+ Project: Restoring and Conserving Meghalaya's Hills Forests through Community Action
PROJECT LOCATION	Khasi Hills, Meghalaya, India
PROJECT SIZE	27,139 ha.
PROJECT DESIGN ORGANIZATION	Community Forestry International (CFI)
PROJECT IMPLEMENTER	Ka Synjuk Ki Hima Arliang Wah Umiam ("FEDERATION")
CHIEF COMMUNITY FACILITATOR	Tambor Lyngdoh
CONTACT INFORMATION	tamborlyngdoh@yahoo.co.in
	Mawphlang-793121, East Khasi Hills District, Mawphlang, India
	Tel: 91-98-63082456 / 91-0364-2567565
TECHNICAL SUPPORT	Technical Advisory Committee (Federation TAC)
	CFI, BioClimate, Rainforest Alliance

TABLE OF CONTENTS

SUMMARY OF PROJECT.....	1
ACRONYMS	3
VERSION CONTROL	3
1. INTRODUCTION	4
2. APPLICABILITY.....	4
3. GEOGRAPHICAL AREA	4
3.1. Climate	6
3.2. Land Cover Types.....	6
3.3. Deforestation and Degradation	6
3.4. Legal Status and Community Rights	8
4. PROJECT ACTIVITIES	9
4.1. Reducing Emissions from Deforestation & Degradation (REDD+).....	9
4.2. Assisted Natural Regeneration (ANR)	10
5. PROJECT PERIOD	11
5.1. Project Timeline	11
6. CARBON POOLS	11
7. BASELINE SCENARIOS	12
7.1. Initial Carbon Stocks.....	12
7.2. REDD+	14
7.3. Assisted Natural Regeneration (ANR)	15
8. PROJECT SCENARIO	16
8.1. REDD+	16
8.2. Assisted Natural Regeneration (ANR)	17
8.3. Leakage.....	18
8.4. Sustainability.....	19
9. CARBON BENEFITS	20
9.1. REDD+	20
9.2. Assisted Natural Regeneration (ANR)	20
9.3. Total Benefits	20
10. MONITORING PLAN.....	21
10.1. REDD+ Monitoring.....	21
10.2. ANR Monitoring.....	23
10.3. Environmental & Biodiversity Indicators.....	24
10.4. Socio-Economic Monitoring	25
10.5. Satellite Monitoring.....	26
10.6. Verification of Targets and Thresholds	26
11. REFERENCES	27
APPENDICES	28
Appendix 1: Mawphlang Pilot Project.....	28
Appendix 2: Biomass Survey	29
Appendix 3: Satellite Image Analysis	31
Appendix 4: REDD+ Project Effectiveness.....	34
Appendix 5: Minimizing the Risks of Non-sustainability	35
Appendix 6: Satellite Image Analysis 2010-2016.....	37

ACRONYMS

ANR	Assisted Natural Regeneration
IGA	Income Generating Activities
LWC	Local Working Conditions
NRM	Natural Resource Management
NTFPs	Non-timber Forest Products
REDD	Reducing Emissions from Deforestation and Forest Degradation
VNRMPs	Village Natural Resource Management Plans

VERSION CONTROL

This version 4.0 of the Technical Specification (November 2018) provides an update to, and follows Version 3.0 (January 2017). The following changes have been made:

- Revised estimates of carbon stocks in dense and open forest, and updated tables to reflect changes to estimates of emissions and removals, as a result of:
 - Adding below-ground woody biomass as an accounted carbon pool
 - Updating volume equations used to estimate individual tree biomass, to use species specific equations when available
 - Applying a more conservative interpretation of mean biomass values by adopting the lower 90% confidence interval
- Revised baseline scenario: correction of a minor error in the interpretation of the satellite analysis. However, the revised values make use of the same analysis as that used for the initial version of the Technical Specification.
- Revised effectiveness of project activities using an analysis of deforestation and degradation observed in an analysis of satellite images from 2010 and 2016. The effectiveness values are applied for the period 2017 to 2021 to give a more conservative estimate of project effectiveness in this period.
- Revised estimated uptake from Assisted Natural Regeneration in 2017 to 2021 to reflect the annual area that will be planted in this period.

1. INTRODUCTION

The Khasi Hills project will slow, halt, and reverse the loss and degradation of forests in North-eastern India. It is India's first community REDD+ project to be certified under an international standard.

Restoration of degraded forests are being achieved by supporting communities in land management and forest regeneration activities in order to yield livelihood benefits. The project supports the development of community natural resource management (NRM) plans for the management of forests and micro-watersheds. Where possible, the project aims to link forest fragments to enhance hydrological and biodiversity services, especially on major and minor riparian arteries of the Umiam River

2. APPLICABILITY

The project represents an innovative approach to community-based forest conservation and restoration that has broad application in the neighboring watersheds in the Khasi hills, as well as more broadly across Meghalaya. The project also seeks to build community institutional capacity to monitor changes in forest cover, hydrological conditions, and biodiversity. The project is located on the traditional forest lands of the Khasi people, which are recognized by the Government of India as community forests under the Sixth Schedule of the Constitution.

This technical specification for reducing emissions from deforestation and forest degradation (REDD+) and assisted natural regeneration (ANR) has been developed for community forests in Meghalaya, India. REDD+ is applicable to dense or open forest under threat of deforestation or degradation. ANR is applicable to open forest. Definitions for dense and open forest are taken from the Indian Forest Survey. Dense forest has canopy cover from 40-100%, while open forest has canopy cover from 10- 40%

3. GEOGRAPHICAL AREA

The REDD+ project is located in the East Khasi Hills District of Meghalaya in Northeast India. The project boundary is the boundary of the Umiam River sub-watershed plus a one-kilometer belt. The project area includes the traditional territories of the nine participating Khasi governments (Hima). The project area is 27,139 hectares. The Umiam sub-watershed is in the Central Plateau Upland region of Meghalaya, India. The altitude of the plateau varies from 150 m to 1,961 m above mean sea level. The plateau has steep regular slopes to the south where Meghalaya borders Bangladesh. The Umiam sub-watershed has rolling uplands, rounded hills and rivers. The River Umiam, which flows through the project area, is a major river in Meghalaya and an important source of water for the capital city of Shillong. Figure 1 shows the Meghalaya Plateau between the Eastern Himalayas and the Arakan Mountains and the locations of the initial 15 community conservation areas.

Figure 1: Meghalaya in Northeast India





Figure 2: Project Area



3.1. Climate

The climate of the Central Plateau Upland region is influenced by its topography and has high seasonal rainfall. There are four seasons: 1) a cool spring from March to April; 2) a hot rainy summer (monsoons) from May to September; 3) a pleasant autumn from October to mid-November and; 4) a cold winter from mid-November to February. The mean maximum temperature ranges between 15°C to 25°C and the mean minimum temperature ranges between 5°C to 18°C.

3.2. Land Cover Types

In the Umiam sub-watershed, there are pine forests, small areas of mixed evergreen cloud forest, barren land, active agricultural land, fallow land and settlements. Forest on community land is mainly pine forest, which is a secondary forest type. Mixed broadleaved and pine forest is found in valleys.

Pure broadleaved forest remnants are confined to gullies, steep slopes and sacred groves. Fragments of mixed evergreen cloud forest remain in the project area as sacred groves. The Mawphlang Sacred Forest is one of the most famous sacred groves in the Khasi Hills (Meghalaya Tourism, 2012). The project area is stratified into four land cover types; 1) dense forest 2) open forest 3) barren or fallow lands and 4) agricultural land (see Table 1).

Table 1: Land Cover Types (Source: Satellite image analysis - see Appendix 3) ¹

LAND COVER	AREA IN 2010 (Ha)
Dense forest	9,270
Open forest	5,947
Barren or fallow	6,330
Agriculture	4,777
Other (shadow/water/no data) ¹	814
Total Area	27,139

3.3. Deforestation and Degradation

The East Khasi Hills have experienced rapid, unplanned deforestation and forest degradation due to social and economic forces. A recent forest survey of India showed that the deforestation rate is 3.6% for dense forest and over 6.8% per year for open forest in the East Khasi Hills District (FSI, 2006) (see Table 2 below).

Table 2: Forest Cover in the East Khasi Hills District: 2001 and 2005. [Source: FSI (2006)]

YEAR	DENSE FOREST	OPEN FOREST	TOTAL
2001	997	1,553	2,550
2005	817	1,019	1,836
Percentage Loss over 5 years	18 %	34 %	28 %
Percentage Loss per year	3.6 %	6.8 %	5.6 %

¹ In the satellite image analysis some areas could not be classified due to water, shadows, or insufficient data. These areas have been grouped into the category called "other" and are treated as non-forest land.

3.3.1. Drivers of Deforestation and Degradation

The key drivers of deforestation and forest degradation in the project area are: 1) forest fires; 2) unsustainable fuel wood collection; 3) charcoal making; 4) stone quarrying; 5) uncontrolled grazing; and 6) agricultural expansion.

With many of the drivers above, it is difficult to estimate the specific contribution of each driver to overall emissions from the project area, due to the limited availability of quantitative data. Firewood consumption, however, lends itself best to such an exercise as virtually all households use firewood with estimated consumption around 10 to 20 kg per household per day. With 4,400 households this means around 15,000 to 30,000 tons of fuelwood are burned each year.

The project estimates that the adoption of fuel-efficient stoves can reduce fuelwood consumption by 30 to 50%, however the project conservatively estimated actual reductions at 15% in the first 5 years and 25% in the second five years. In addition, NRM plans will include rotational fuelwood harvesting and the establishment of fast growing plantations that should increase fuelwood supplies, reducing the rate of forest degradation and accelerating natural regeneration by reducing pressure on the natural forests. Plantations may generate around 4,000 to 6,000 tons of fuelwood a year once they are productive (year 5) which would meet approximately 30 to 40% of the demand in the project area.

The Federation will be monitoring changes in forest conditions and the drivers of forest loss and degradation. Feedback on forest loss will be communicated by LWC members to the Federation on an annual basis. In addition, every five years, updated satellite images of the area will be analyzed to identify where forest loss is occurring. Based on this information the Federation will identify the causes and appropriate mitigation measures. Risk from natural hazards appears low at the present time. This upland region is not in a flood zone, nor is it subject to landslides or frequent earthquakes.

3.3.2. Forest Fires

Fires occur during dry months when the forest floor is covered with a thick layer of dry leaves and needles. In recent years, ground fires are estimated to burn approximately 25% of the project area each year. Fires are often set by discarded cigarettes, children playing with matches and escaping fires from agricultural burning. CFI's earlier pilot project demonstrated that community awareness-raising with community imposed prohibitions on smoking and carrying matches into the forest have significantly reduced the incidence of fire. Building fire-lines and hiring village firewatchers also contributed to reductions in ground fires. In addition, the establishment of fines for those who cause fires also creates an incentive to be careful. Incidence of fire will be monitored by the LWC as burn areas are highly visible. Rewards to communities that prevent fire may be given at the end of the fire season. Training in fire safety and control is also important as communities may use fire to establish fire-lines (sanding) as well as for agricultural clearing.

3.3.3. Unsustainable Firewood Collection

Over 99% of the rural community uses firewood as their sole source of fuel. Being situated in a relatively cold region, firewood consumption per household in the area is high, averaging 10 to 20 kg per household per day. Firewood is collected from nearby forests. If dead trees are not available, people resort to felling live trees and saplings. While some villages have regulations guiding

fuelwood collection, many do not or these systems have broken down. The establishment of an NRM (plan vivo) planning process will help communities re-establish sustainable firewood production systems.

3.3.4. Charcoal Production

There is a significant demand for charcoal in Meghalaya. Charcoal is used by iron-ore smelting industries and it is also used for heating homes and offices in urban centers such as the city of Shillong. Charcoal making and its purchase by industries is illegal in Meghalaya. Charcoal making is concentrated in a few villages with limited alternative income generating opportunities.

3.3.5. Stone Quarrying

There is a large demand for stone, sand and gravel for construction in Shillong city. Many stone quarries exist in the project area. Quarries are usually on steep slopes and they lead to erosion and landslides. Hima governments will be asked to place a moratorium on leasing land for quarries and not extend existing leases wherever possible.

3.3.6. Uncontrolled Grazing

The rural communities allow cattle, goats and sheep to graze in nearby forest areas. Grazing causes forest degradation as young seedlings and saplings are grazed or trampled. Grazing animals are reported to have little economic value with communities often eager to switch to stall feeding and higher quality livestock.

3.3.7. Agricultural Expansion

Communities or clans own most of the forests in the project area. However, when community and clan forests are privatized they are often permanently cleared for agriculture. Forest clearance is also practiced for extensive and shifting agriculture (*jhum*) on steep slopes.

Agricultural expansion is taking place in several Hima in the southern part of the project area where businessmen are providing loans to families to clear forests and plant broom grass for markets in other parts of India. Slowing and halting this process will require consultations with farmers involved in this activity to discuss alternative agricultural and other economic activities which could be supported both through the project as well as under Government of India schemes and projects.

3.4. Legal Status and Community Rights

There are no legally designated or protected conservation areas within, overlapping or adjacent to the project area. Communities own their land through a legally recognised land-tenure system. In this system, *Dorbars* are the administrative heads of territorial units, and decisions regarding community land are made by consensus by male community members over 21 years of age.

4. PROJECT ACTIVITIES

REDD+ and ANR are the Plan Vivo project interventions covered in this technical specification. REDD+ is the protection of dense or open forest threatened by deforestation and forest degradation. ANR is the protection, management, and regeneration of open forest. In addition to REDD+ and ANR interventions, other income-generating activities (IGAs) are designed to improve local livelihoods. IGAs have been designed by the communities and are facilitated by the project team.

4.1. Reducing Emissions from Deforestation & Degradation (REDD+)

REDD+ intervention addresses the key drivers of deforestation and forest degradation in the project area. It consists of the following activities: 1) forest fire control, 2) sustainable firewood plantations, 3) reducing uncontrolled grazing and 4) agricultural containment.

4.1.1. Forest Fire Control

Damage from forest fires will be reduced through fire prevention and early fire detection. Activities to control forest fires include:

- Creating firebreaks around forests
- Appointing firewatchers to detect and extinguish fires in the dry season
- Community fire awareness programmes to improve fire safety

4.1.2. Sustainable Firewood

Sustainable firewood plantations will be established close to settlements and firewood gathering will be organized around a rotational system of harvesting with guidelines for fuel collection during years 1 to 5 as the fuelwood plantations grow and mature. Fuelwood collection areas are associated with specific villages, so that there is limited likelihood of displacement or leakage from other communities outside the project area. With the project, fuelwood access is more regulated based on emerging NRM plans. Project woodlots will take 4-5 years before annual harvesting of coppice shoots takes place. Of the 5,947 ha of forest in the project area, woodlot plantations will likely cover approximately 300 ha (5 ha for each village), depending on funding availability.

4.1.3. Reduce Uncontrolled Grazing

Through animal exchange programs, communities will be encouraged to replace cattle with stall-fed livestock such as pigs and broiler chickens. The Mawphlang Pilot Project demonstrated that participating families were able to transition from open forest grazing with low value goats and cows to stall fed pigs, reducing pressure on the forests while generating additional income from pig sales.

4.1.4. Sustainable Farming Systems

The project will support the adoption of sustainable agricultural practices. Sustainable agriculture refers to farming systems that are likely to be practiced for extended periods without damage to forests and soils. This would include organic vegetable cultivation and orchards, stall fed livestock, and aquaculture. Unsustainable systems such as broom grass, pineapples requiring the clearing of vegetation on steep slopes, and valley bottom potatoes requiring high use of chemical fertilizers and pesticides will be phased out where possible. The project is building partnerships with the Indian Council for Agricultural Research that provides training and materials for exploring new agricultural practices. Project funded micro-finance groups provide capital for small farmers to adopt sustainable farming practices.

4.1.5. Alternatives to Charcoal Making

Charcoal making is concentrated in two of the 10 Project Hima. In those areas, meetings are being planned with charcoal-making households to identify alternative livelihood activities including pig and poultry raising. Funds will be allocated to provide support to these families to help them transition their household economy.

The core project strategy begins with a community dialogue followed by an agreement on the part of all member households to attempt to reduce the impact of drivers of deforestation activities and build mitigation activities into their NRM plan (Plan Vivo). As mentioned above this approach has been implemented with considerable success in two villages in the project area from 2005 to 2009. The project has a successful approach to replacing low value cows and goats with stall fed chicken and pigs (see PDD) reducing grazing pressures.

Fire control efforts of the community were very successful through 5 fire seasons. Agricultural expansion is most threatening where forests are cleared for cash crops, especially broom grass. Areas where this is occurring have been identified and targeted discussions with practitioners are planned to find more sustainable crops outside the forests. Reducing charcoal making will again target the charcoal making households to help them find alternatives. Involving female members in micro-finance self-help groups and providing technical training and low interest loans to establish piggeries and poultry operations.

4.2. Assisted Natural Regeneration (ANR)

ANR activities will take place in open forest. There are two ANR phases. The initial phase of “advanced closure” involves “closing” the area to fire, grazing, and firewood collection. The second “ANR treatment” phase involves weeding, thinning, and enrichment plating. No exotic species will be used in the ANR areas. Some limited gap filling and enrichment planting will take place using native Khasi pine saplings (*Pinus khasiana*) as well as oak (*Quercus griffithi*), chestnut (*Castanopsis purpurella*) and myrica (*Myricaesculenta*).

A long term goal of the project is to improve the soil fertility, soil moisture, biomass, and species diversity of the open forests through ANR treatment. Past experience from the Mawphlang pilot project (2005-2009) indicated that with protection through advanced closure, forest regrowth was quite rapid. Open forests tend to be dominated by pioneering Khasi pine seedlings that grow quickly in many sites once grazing, hacking and fire pressures are removed. Over time, a growing number of native broadleaved and evergreen species of shrubs and trees emerge creating more diverse forest ecology. In sites, with no seed sources, enrichment planting of native oaks and

chestnuts will be encouraged to facilitate this process.

ANR advance closure will be implemented in 25 of the open forest in the first implementation phase (2012-2016), expanding to 30% of the area in the second implementation phase (2017-2021) (see Table 3).

Table 3: ANR Area

ANR TREATMENT TYPE	IMPLEMENTATION PHASE 1 2012-2016 (Ha)	IMPLEMENTATION PHASE 2 2017-2021 (Ha)	TOTAL 2012-2021 (Ha)
ANR advance closure	1,401	408	1,809
ANR treatment	500	500	1,000

5. PROJECT PERIOD

The initial project period is 5 years. At the end of this period (2016) and following verification, the project was extended and the baseline and technical specifications duly updated.

The total project period is 30 years. For the carbon benefit calculation, the project period is divided into three five-year crediting periods.

5.1. Project Timeline

From 2005 to 2009, CFI organized REDD+ and IGA pilot activities in two communities in Mawphlang (Appendix 1). Following the success of the Mawphlang pilot project, the design process for the Khasi Hills REDD+ project took place in 2010-2011. In 2011-2012, early REDD+ activities including institution building, awareness campaigns, field activity development, and the design of monitoring systems began. The first implementation phase of the project took place from 2012 to 2016. At the end of the first implementation phase, the Technical Specifications were revised and the expected climate benefits have been updated prior to the second implementation phase of the project from 2017 to 2021.

6. CARBON POOLS

Above- and below-ground tree biomass are the carbon pool used to calculate carbon benefits for both REDD+ and ANR (see Table 4). Other carbon pools are omitted for three reasons: simplicity, cost of measurement and conservativeness.

Including only tree biomass leads to simple and less resource- intensive monitoring, measurement, and analysis. The resulting carbon benefit estimate is also conservative as the storage and sequestration in soil and, deadwood and litter, are not being claimed as credits by the Project. Consequently, this represents a buffer that may help reduce project risk.

Explanations for carbon pool selection are:

- Above- and below-ground tree biomass comprise the main carbon pools - these are included
- Biomass stored in leaf litter and dead wood will increase as a result of tree-planting activities, but is unlikely to be a large proportion of the total carbon and is therefore

excluded

- Non-tree vegetation is unlikely to be a large proportion of the total carbon stock and is excluded
- Soil carbon is expected to increase but the cost of measuring it is high, so it is excluded
- Dead wood is likely to increase during forest conservation, but this is not included to allow a conservative estimate of carbon benefit

Table 4: Carbon Pools

CARBON POOL	LIKELY IMPACT ON CARBON STOCK	MEASUREMENT LIMITATIONS	DECISION
Above-ground woody biomass	Increase	Minimal	Include
Below-ground woody biomass	Increase	Minimal	Include
Non-tree biomass	Small increase	Time-consuming	Exclude
Dead wood	Increase	Minimal	Exclude
Leaf litter	Small increase	Time-consuming	Exclude
Soil	Increase	Expensive	Exclude

7. BASELINE SCENARIOS

7.1. Initial Carbon Stocks

Initial carbon stocks in the project area were determined by carrying out a biomass survey and a satellite image analysis (see Table 5).

Table 5: Carbon Stock in 2010

LAND USE	AREA (Ha)	TOTAL CARBON STOCK (tC)	TOTAL CARBON STOCK (tCO ₂ e)
Dense forest	9,270	878,193	3,220,042
Open forest	5,947	71,761	263,123
Non-forest	11,921	0	0
Total	27,139	949,954	3,483,165

Source: Biomass Survey (see Appendix 2)

Biomass Survey

The project team carried out a biomass survey of 21 plots in dense forest and 19 plots in open forest (Appendix 2) in 2010 to assess initial carbon stock. Dense and open forest areas were identified on a land cover stratification map based on remote sensing data from the Forest Survey of India (2004), contour maps and path network maps. Most of the forestland is relatively inaccessible, far from roads or tracks or on steep slopes and plateaus cut by gullies and cliffs. For this reason, sample plots were selected randomly along transects that follow the existing local path network running east- west and north-south. Dense forest plots were 10 square meters (0.02 ha), and open forest plots were 20 square meters (0.04 ha). In each plot, the tree species and diameter at breast height (DBH) were recorded as well as top heights of three trees at the lower, middle, and upper canopy (Table 6).

To calculate biomass from sample plot measurements for dense forest plots, species-specific volume equations (FSI 1996; Table 7) were used to estimate stem volume of individual trees. The Forest Survey of India – based on measurements of the tree dimensions during past fellings of thousands of trees over two decades – developed these equations. If species-specific equations were not available a generic equation for north-east Indian tree species was used. Stem volume was converted to stem biomass by multiplying the volume estimate by species-specific wood density values for trees in India from the Global Wood Density Database (Zanne 2009). If species-specific wood density values were not available a values of 0.652 g/cm³ was applied, which is the average of all Indian species in the Global Wood Density Database.

A biomass expansion factor (BEF) was then applied to convert stem biomass estimates to estimates of whole tree biomass was applied. Biomass expansion factors recommended by Brown (1997) were applied:

- When inventoried biomass was >190 t/ha a BEF of 1.74 was applied;
- When inventories biomass as <190t/ha a BEF = EXP(3.213-0.506*LN(BV)), was applied where BV=inventoriedvolume;
- For plots dominated by pines a BEF of 1.3 was applied.

Below-ground biomass was estimated by assuming a root:shoot ratio of 0.15 for all species. FSI (1996) reports a range of root-shoot ratios, with values up to 0.32. Other studies in Punjab (e.g. Rawat et al. 2015) report lower values however, ranging from 0.15 to 0.19 depending on tree age. To avoid overestimating below ground biomass, the most conservative value from the literature was selected.

Since there is some uncertainty in estimated biomass from tree inventories, related to the variation in biomass between sample plots in the same forest type, the lower 90% confidence interval of mean values was adopted to estimate biomass for each forest type. Estimated biomass was therefore 12.1 tC/ha for open forest and 94.7 tC/ha for dense forest. It was assumed that carbon stock for barren or fallow land and agricultural land was zero. Inventories of sample plots will be taken again at the end of the initial Project period in 2015.

Table 6: Biomass Survey Values.

Open Forest		Dense Forest	
Plot No.	tC/ha	Plot No.	tC/ha
3	16.9	1	170.5
6	3.3	2	97.1
7	10.3	4	86.7
8	6.3	5	57.9
12	19.5	9	160.2
13	8.5	10	108.9
15	28.4	11	100.1
16	54.1	14	144.8
17	44.3	21	87.4
18	12.7	22	151.7
19	8.9	24	125.9
20	5.9	25	21.1
23	8.2	28	75.8
26	24.8	29	140.2
27	35.0	30	209.9
31	10.5	33	63.0
36	11.9	34	119.8

39	22.2	35	53.8
40	4.4	37	213.2
		32	63.5
		38	142.3
Mean	17.7	17.6	114.0
90% CI	5.6	5.3	19.3

Source: Biomass Survey (see Appendix 2)

Table 7: Local volume equations for different species for Meghalaya state.

SPECIES	VOLUME EQUATION *	NOTE
<i>Castanopsis hystrix</i>	$V=0.13937-0.35988VD+6.81318D^2$	
<i>Castanopsis indica</i>	$V=0.22234+4.90695D+1.5124VD$	
<i>Engelhardtia spicata</i>	$\text{Log}_e V=2.47635+2.51046 \text{ Log}_e D$	
<i>Pinus kesiya</i> **	$V=0.0232-0.011613D+0.0011549D^2$	(diameter in cm)
<i>Quercus fenestrata</i>	$V/D^2=0.000295/D^2-0.0079835/D+0.00086$	(diameter in cm)
<i>Quercus glauca</i>	$V/D^2=0.000295/D^2-0.0079835/D+0.00086$	(diameter in cm)
<i>Quercus griffithii</i>	$V/D^2=0.000295/D^2-0.0079835/D+0.00086$	(diameter in cm)
<i>Rhododendron arboreum</i>	$V=0.08934+0.70730D+2.13941D^2$	
<i>Schima wallichii</i>	$V=0.27609-3.68443D+15.866870D^2$	
<i>Symplocos theaefolia</i>	$V=0.03754+0.000587D^2$	(diameter in cm)
<i>Others</i>	$V=0.11079-1.81103D+11.4132D^2+0.38528D^3$	

Source: FSI 1996. Note: Equations selected were those derived from measurements of trees in closest proximity to the project site.

* V=Volume in m^3 ; D = Diameter in m (unless specified otherwise)

** If the dbh of the pine trees are <10 cm, the generic volume equation was applied, as the species specific equation was not intended for use on trees <10 cm diameter.

7.2. REDD+

A SPOT satellite image analysis was carried out to determine the land-use types and areas present in 2006 and 2010 (Table 8) as well as the recent rates of forest degradation and deforestation (Table 9). See Appendix 3 for a detailed description of the satellite image analysis.

Table 8: Land Use and Land Cover in 2006 and 2010.

LAND USE	2006 (Ha)	2010 (Ha)
Dense forest	10,446	9,270
Open forest	5,908	5,947
Barren or fallow	5,794	6,330
Agriculture	3,179	4,777
Other (shadow/water/no data) ¹	1,812	814
Total Area	27,139	27,139

Source: Satellite Image Analysis (see Appendix 3)

From 2006 to 2010, dense forest changed to non-forest land at a rate of 2.7% per year; dense forest changed to open forest at a rate of 4.5% per year; and open forest changed to non-forest at a rate of 6.4% per year (Table 9).

Table 9: Forest Degradation and Deforestation from 2006 to 2010.

LAND USE CHANGE (2006 TO 2010)		AREA (HA)	CHANGE PER YEAR
Dense forest changed to non-forest		1,136	2.7%
Dense forest changed to open forest		1,860	4.5%
Open forest changed to non-forest		1,510	6.4%

Source: Satellite Image Analysis (see Appendix 3)

For the REDD+ baseline scenario we assume recent rates of deforestation and degradation would continue over the next ten years in the absence of project activities, resulting in 1,845 ha of dense forest being changed to non-forest; 3,700 ha of open forest changed to non-forest; and 3,021 ha of dense forest being changed to open forest in the project area. This results in 1,720,050 tCO₂ emissions (Table 10).

Table 10: REDD+ Baseline

Year	Dense forest changed to non-forest (ha)	Open forest changed to non-forest (ha)	Dense forest changed to open forest (ha)	C stock change from conversion of dense forest to non-forest (tC)	C stock change from conversion of open forest to non-forest (tC)	C stock change from conversion of dense forest to open forest (tC)	Total C stock change (tC)	Total emissions (tCO ₂)*
2012	252	380	413	23,875	4,585	34119	62,578	229,453
2013	234	382	383	22,163	4,610	31672	58,445	214,298
2014	217	382	356	20,574	4,611	29401	54,585	200,146
2015	202	380	330	19,098	4,590	27293	50,981	186,931
2016	187	377	306	17,729	4,552	25335	47,616	174,592
2017	174	373	284	16,458	4,497	23519	44,473	163,069
2018	161	367	264	15,277	4,429	21832	41,539	152,309
2019	150	360	245	14,182	4,350	20267	38,798	142,260
2020	139	353	228	13,165	4,261	18813	36,239	132,877
2021	129	345	211	12,221	4,164	17464	33,849	124,114
Total	1,845	3,700	3,021	174,742	44,648	249,715	469,104	1,720,050

Source: See tables 5 and 9.

* Assuming all reductions in C stock result in instantaneous emission of CO₂

7.3. Assisted Natural Regeneration (ANR)

ANR activities will be implemented in open forest areas. In the absence of project activities, it is assumed that open forests would continue to degrade due to periodic forest fires, unsustainable fuelwood extraction, agricultural expansion and grazing, gradually losing biomass, rootstock, and top soil. Typically, under the without project scenario, new shoots are hacked for firewood, seedlings are trampled by cattle and goats, and ground fires retard or destroy seedlings and saplings. This pattern has been observed throughout the project area leading up to the initiation of the project. Carbon stocks in open forest are therefore expected to decline in absence of project interventions, but the rate of decline is not known. We therefore adopt the conservative assumption that carbon stocks in open forests would remain constant at 12.1 tC/ha in the absence of project interventions.

8. PROJECT SCENARIO

8.1. REDD+

During the ten-year project period, there are two implementation phases. In the first implementation phase (2012-2016), activities were started, and in the second implementation phase (2017-2021), activities will be intensified in terms of participation.

At the start of the project it was estimated that the overall rate of deforestation and forest degradation without the project would be reduced by 33% at the start of the first implementation phase, increasing to 57% by the end of the second implementation phase (see Table 11 and Appendix 4). Under the original REDD+ Project scenario, emissions would be 971,548 tCO₂e over 10 years (see Table 12). The estimated 33% decrease at the start of the first five years was based on impacts achieved between 2005 and 2010 in the original pilot project area where ground fires were dramatically reduced, as were grazing and fuelwood collection pressures. The expected effectiveness of community mitigation measures was due to the consensus decision taken by the indigenous government (Hima Mawphlang) and the participating village durbar meetings and discussions. This REDD+ project has adopted the same approach and is being developed at the request of 10 neighboring indigenous governments that have seen the results of the pilot activities.

Table 11: Original REDD+ Project Scenario (2010).

Year	Reduction in deforestation and degradation % to non-forest (ha)	Dense forest changed to non-forest (ha)	Open forest changed to non-forest (ha)	Dense forest changed to open forest (ha)	C stock change (dense forest to non-forest) (tC)	C stock change (open forest to non-forest) (tC)	C stock change (dense forest to open forest) (tC)	Total C stock change (t C)	Total emissions (t CO ₂)*
2012	33%	169	255	277	15,996	3,072	22,859	41,927	153,734
2013	36%	150	245	245	14,184	2,950	20,270	37,405	137,151
2014	38%	135	237	221	12,756	2,859	18,229	33,843	124,091
2015	41%	119	224	195	11,268	2,708	16,103	30,079	110,290
2016	44%	105	211	172	9,928	2,549	14,188	26,665	97,771
2017	46%	94	201	154	8,887	2,428	12,700	24,016	88,057
2018	49%	82	187	135	7,791	2,259	11,134	21,185	77,677
2019	52%	72	173	118	6,807	2,088	9,728	18,623	68,285
2020	54%	64	162	105	6,056	1,960	8,654	16,670	61,123
2021	57%	55	148	91	5,255	1,791	7,510	14,555	53,369
Total		1,044	2,044	1,710	98,929	24,664	141,375	264,968	971,548

Source: Appendix 4, Tables 5 and 10.

* Assuming all reductions in C stock result in instantaneous emission of CO₂

At the end of the first implementation phase, in 2016, analysis of remote sensing data was carried out to describe the change in land cover that occurred in the period from 2010 to 2016 (see Appendix 6). The estimated effectiveness of the project was revised for the second implementation period (2017 to 2021). The results suggest that, relative to the baseline scenario, project activities in the first project implementation period reduced deforestation of dense forest by 20.2%, reduced deforestation of open forest by 28.5% and reduced degradation of dense forest to open forest by 35.0% (see table 12).

Table 12: Forest Degradation and Deforestation from 2010 to 2016.

LAND USE CHANGE (2010 TO 2016)	TOTAL AREA (Ha)	ANNUAL AREA (Ha)	CHANGE PER YEAR	EFFECTIVENESS RELATIVE TO BASELINE
Dense forest changed to non-forest	1,328	221.4	2.2%	20.2%
Dense forest changed to open forest	1,770	295.1	2.9%	35.0%
Open forest changed to non- forest	1,029	171.4	4.6%	28.5%

Source: Satellite Image Analysis (see Appendix 6), and Table 9.

For the revised REDD+ project scenario, for the second project implementation period (2017 to 2021), it is assumed that the project activities will result in the same level of effectiveness achieved during period from 2010 to 2016 (see Table 12). This is expected to provide a conservative estimate of project effectiveness for two reasons. Firstly because the estimated effectiveness from 2010 to 2016 includes two years during which project activities were not fully operational, and secondly because this does not factor in the expected increase in effectiveness as the project progresses that was included in the original project scenario.

The revised estimate of emission reductions from REDD+ achieved during the first phase of the project, and expected during the second phase of the project, are summarized in Table 13. The revised project scenario has higher total emissions than expected with the effectiveness assumed in the original project scenario.

Table 13: Revised REDD+ Project Scenario (2016)

Year	Dense forest changed to non- forest (ha)	Open forest changed to non- forest (ha)	Dense forest changed to open forest (ha)	C stock change (dense forest to non-forest) (tC)	C stock change (open forest to non-forest) (tC)	C stock change (dense forest to open forest) (tC)	Total C stock change (t C)	Total emissions (tCO ₂)*
2012	252	380	413	23,875	4,585	34,119	62,578	229,453
2013	187	273	249	17,697	3,294	20,580	41,571	152,426
2014	173	273	231	16,428	3,294	19,105	38,827	142,364
2015	161	272	215	15,250	3,280	17,735	36,264	132,968
2016	149	270	199	14,156	3,252	16,463	33,871	124,194
2017	139	266	185	13,141	3,213	15,282	31,637	116,001
2018	129	262	172	12,199	3,165	14,186	29,550	108,349
2019	120	258	159	11,324	3,108	13,169	27,601	101,204
2020	111	252	148	10,512	3,044	12,225	25,781	94,531
2021	103	247	137	9,758	2,975	11,348	24,082	88,299
Total	1,524	2,752	2,107	144,340	33,210	174,212	351,761	1,289,791

8.2. Assisted Natural Regeneration (ANR)

Assisted natural regeneration (ANR) will take place in open forest. ANR activities begin with “advance closure” to protect the area from fire and grazing and to allow the trees to regenerate. Following advance closure, some areas also receive “ANR treatment” which is weeding, thinning and enrichment planting.

All ANR activities have the same initial advance closure stage and therefore the same carbon benefit. Based on our observation of rapid forest regeneration in the Mawphlang project (Appendix 1), we assume that open forests can regenerate into dense forest in 30 years. Assuming open forests can regenerate into dense forests in 30 years, the average annual carbon sequestered would be approximately 1.95 tC/ha/yr; however, we make the conservative assumption that open forest with ANR will sequester carbon at a rate of 1 tC/ha/yr for the first 10 years and a rate of 1.5 tC/ha/yr for the following 20 years.

The ANR sequestration rates estimated for the project compare well with findings from studies of similar open pine forests. Open pine forests can sequester carbon at a rate between 1.07 and 1.6 tC/ha/yr (Table 14). The related studies from central Nepal are based on degraded Chir pine forests that are very similar to the khasi pine (*Pinus khasiana*) that dominates the open forest landscape in the project area. Further, elevation is similar, though rainfall in the project area is considerably higher than western Nepal, suggesting that growth in the project area may be more rapid.

Table 14: Carbon Sequestration in Open Pine Forests

REFERENCE	OPEN PINE FOREST (tC/Ha/Yr)
Shrestha, R. (2010)	(1.6 pine + 1.37 poor condition)/2 = 1.5
Baral et al, (2009)	1.35 (pine)
Jina et al, (2008)	1.07 to 1.27 (degraded pine)

Between 2013 and 2016 the project worked with communities to bring 1,401 hectares under ANR with plans to bring an additional 408 hectares under ANR between 2017 and 2021. The climate benefits from ANR achieved during the first phase of the project, and expected during the second phase, are summarized in Table 15.

Table 15: Estimated Carbon uptake achieved during Phase 1 and expected during Phase 2.

Year	Area Planted (Ha)	Cumulative Area Planted (Ha)	Carbon Uptake (tC)	Emissions Reductions (tCO ₂)
2012	0	0	0	0
2013	0	0	0	0
2014	392	392	392	1,439
2015	501	893	893	3,277
2016	500	1,393	1,393	5,112
2017	8	1,401	1,401	5,142
2018	100	1,501	1,501	5,509
2019	100	1,601	1,601	5,876
2020	100	1,701	1,701	6,243
2021	100	1,801	1,801	6,610
Total	1,801	10,683	10,683	39,308

In grey: actual numbers until 2018

8.3. Leakage

For each risk of leakage, the project includes leakage mitigation measures for both REDD+ and ANR (See Table 16). With leakage mitigation measures in place, activities causing emissions are unlikely to be displaced outside the project area. Therefore, we assume the risk of leakage risk to be low and have applied a 5% leakage deduction to the overall benefit calculations for both REDD+ and ANR.

Table 16: Leakage Mitigation Measures

Drivers of Mitigation	Activity	Mitigation Measures
Firewood Collection	REDD+, ANR	Village Natural Resource Management Plans (VNRMPs) will be designed to ensure that firewood requirements are met from community land. VNRMPs will include the establishment of plantations close to villages to supply firewood. This wood will be harvested sustainably using rotational harvesting systems.
Charcoal making	REDD+	Charcoal-making is a driver of deforestation and forest degradation in one of the nine Himas in the project area, Nonglwai Hima. Charcoal making and its purchase by industries are illegal in Meghalaya. Assistance from administrative authorities will be obtained to help to check illicit movement of charcoal to ferro-alloy factories in and around the project area.
Agricultural expansion	REDD+, ANR	The project will introduce sustainable agricultural practices to replace unsustainable swidden farming. This will lead to agricultural containment in the project area, and agricultural expansion will not be displaced outside the project area.
Grazing in forest	REDD+, ANR	Cattle and goats will be exchanged for stall-fed livestock through an animal exchange program. This will reduce grazing in the project area and will not increase the risk of grazing outside the project area.

8.4. Sustainability

REDD+ and ANR activities are designed to be sustainable and to supply benefits after the project period. Firstly, the project team is working to reduce financial, management, and technical risks. Secondly, political, social, land ownership, and opportunity cost risks will be addressed. Thirdly, the risks of fire will be minimized. Please see appendix 5 for a detailed analysis. The risk table attempts to quantify the risk for a range of risk factors including socio-political, institutional, financial, and natural events.

The formula is based on giving a score to the likelihood the risk factor will occur (.05 = unlikely, and .1 = likely) multiplied times the severity of potential impact to the project (1= low, 2= medium and 3= high). This provides a composite score that would suggest a buffer of 20%. Overall the project is comparatively low risk in the South Asia context due to very strong tenure security, active and democratic indigenous governments, high literacy in the project communities, and a strong local commitment to restoring forests in the watershed.

8.4.1. Risk Buffer

The risk buffer is a proportion of carbon benefits that are not sold. It is based on the risk of non-sustainability of the project. We estimate that a 20% risk buffer is appropriate for project activities where Plan Vivo certificates are sold ex-post and in accordance with the Plan Vivo guideline for REDD+ projects. The project design relies on a conservative estimate of carbon stocks and benefits in order to reduce the risks of over-estimating carbon credits generated by the project. Potential carbon offsets from below ground biomass, litter and deadwood are also not included and can be viewed as an additional risk buffer.

9. CARBON BENEFITS

9.1. REDD+

Over 10 years, the expected REDD+ net benefit expected is 408,745 tCO₂ (See Table 17).

Table 17: REDD+ Emissions Reductions

Year	Baseline scenario emissions (tCO ₂)	Project scenario emissions (tCO ₂)	Leakage (tCO ₂)	Emissions Reductions (tCO ₂)
2012	229,453	229,453	0	0
2013	214,298	152,426	3,094	58,778
2014	200,146	142,364	2,889	54,893
2015	186,931	132,968	2,698	51,265
2016	174,592	124,194	2,520	47,878
2017	163,069	116,001	2,353	44,715
2018	152,309	108,349	2,198	41,761
2019	142,260	101,204	2,053	39,004
2020	132,877	94,531	1,917	36,429
2021	124,114	88,299	1,791	34,024
Total	1,720,050	1,289,791	21,513	408,745

9.2. Assisted Natural Regeneration (ANR)

Over 10 years, the expected ANR benefit is 37,247 tCO₂e (see Table 18).

Table 18: ANR Emissions Reductions

Year	Baseline scenario emissions (tCO ₂)	Project scenario emissions (tCO ₂)	Leakage (tCO ₂)	Emissions Reductions (tCO ₂)
2012	0	0	0	0
2013	0	0	0	0
2014	0	-1,439	72	1,367
2015	0	-3,277	164	3,113
2016	0	-5,112	256	4,856
2017	0	-5,142	257	4,885
2018	0	-5,509	275	5,234
2019	0	-5,876	294	5,582
2020	0	-6,243	312	5,931
2021	0	-6,610	331	6,279
Total	0	-39,308	1,961	37,247

9.3. Total Benefits

Table 19 below shows the projected carbon benefits for the first ten years of the project that are estimated to result from all planned Project activities. These estimates have been reviewed in 2016 during the first 5-year verification.

Table 19 shows the annual Project benefit from both REDD+ and ANR over the whole 10-year period. The final column shows the total benefits (per year) with a 20% risk buffer subtracted. The table also reflects the additional 500 ha treated under ANR each year beginning in 2014 with an average sequestration rate of 1tC per ha per year.

Table 19: Total project carbon benefits

Year	Net REDD+ benefit (tCO ₂)	Net ANR benefit (tCO ₂)	Overall project benefit (tCO ₂)	20% Buffer (tCO ₂)	Net Total (minus buffer) (tCO ₂)
2012	0	0	0	0	0
2013	58,778	0	58,778	11,756	47,022
2014	54,893	1,367	56,260	11,252	45,008
2015	51,265	3,113	54,378	10,876	43,502
2016	47,878	4,856	52,734	10,547	42,187
2017	44,715	4,885	49,600	9,920	39,680
2018	41,761	5,234	46,995	9,399	37,596
2019	39,004	5,582	44,586	8,917	35,669
2020	36,429	5,931	42,360	8,472	33,888
2021	34,024	6,279	40,303	8,061	32,242
Total	408,747	37,247	445,994	89,200	356,794

10. MONITORING PLAN

The Project has developed a comprehensive monitoring plan based on the requirements of the Plan Vivo Standard (2013). This plan will enable the Project to monitor performance (assessed by achievement of annual targets and five- year goals), validate assumptions used for calculating the carbon benefits and ensure community involvement. The monitoring plan also includes monitoring of indicators to assess the effectiveness of Project activities to mitigate the key drivers of deforestation and forest degradation and of indicators to assess the socio-economic impacts and environmental impacts of the Project to ensure that these aspects of the Plan Vivo Standard are met. The monitoring Plan is summarized in the following three tables including Table 20: Ecosystems Service Benefit Indicators, Table 21: Socio-Economic Monitoring Indicators, and Table 22: Environmental and Biodiversity Monitoring Indicators.

Baselines were established at the start of the Project in 2011 covering (a) forest cover (b) carbon stock and (c) socio-economic indicators. As a REDD+ project, annual monitoring (and reporting) is largely based on monitoring of activities supported by the project with impact monitoring taking place every 5 years and with the resulting information being used to revise this technical specification.

Indicators measured and recorded annually (see Tables 20, 21, 22 below) will be submitted in the Project Annual Reports submitted to Plan Vivo. Results from five-year indicators will be used during preparation of project verification reports.

10.1. REDD+ Monitoring

The primary methodology used to monitor changes in forest cover is an analysis of a time series of satellite images of the project area. For the baseline, SPOT images from 2006 and 2010 were used to determine that the rate of deforestation was 2.7% per annum. For forest areas that have moved from the dense forest category (40% canopy closure or more) to non-forest, the rate of degradation was 0.1%. For forest areas that have moved from the dense forest category to open forest (10 to 40% canopy closure). Actual changes in forest cover were determined at the end of 2016 through the analysis of satellite image done in 2017 and will be reassessed every 5 years (i.e. 2021, 2026, 2031, etc.).

Data from a biomass survey of sample plots and photo monitoring is used to assess the actual carbon stock. The annual biomass survey includes 20 permanent sample plots that were surveyed for the baseline in April 2011 and will be used for the long term monitoring of carbon stock changes in the dense forests that comprise the REDD+ project area. This method allows an annual assessment of changes in carbon stock for dense forests to be made.

It is estimated that the rate of forest loss will gradually be reduced over the Project period from an initial 33% reduction in the deforestation/degradation rate to a 57% reduction after the end of the first 10-year crediting period compared with the baseline (without Project) scenario. By 2025, it is projected that forest cover will stabilize and begin to expand as open forests recover. As actual rates are monitored, the estimated project benefits will be recalculated and technical specifications revised accordingly.

The Federation has identified on current SPOT images 'hot spots' where dense forest loss is occurring and is meeting with local communities to discuss how to reduce forest loss. Specific drivers are associated with certain 'hot spots' and special attention will be given to monitoring forest clearing for broom grass cultivation as well as charcoal making with discussions with households participating in these activities to find alternative income generating activities. Each Local Working Committee will report annually on forest losses due to specific drivers of deforestation such as broom grass cultivation, charcoal making and others.

In addition to the analysis of remotely sensed data to monitor forest cover, the Project will conduct annual field-level inventories of 60 forest plots to assess changes in biomass and carbon stock. The measurements are conducted at the end of each calendar year. The forest plot sample includes 20 dense forest plots (10m x 10m), 20 open forest plots (20m x 20m), and 20 plots under Assisted Natural Regeneration (ANR) (20m x 20m). The data is collected in November each year and analyzed to assess changes in biomass. The plot locations are marked with paint and identified using GPS coordinates. This will include both the dense forest plots and the open forest/ANR plots. Resources required for monitoring include a forestry professional guide, the community facilitator team that works for the Federation, and members of the LWC who are trained in forest inventory techniques. Equipment includes plot and tree measuring tapes, clipboards and data collection forms, cameras, GPS units, plot lines, and paint. The data will be analyzed by the Federation and the project's REDD+ Technical Support Unit (RTSU) using and EXCEL and ACCESS data base system.

Annually, at the end of the rainy season, monitoring photo will be taken from a known fixed point in the plot. The project has established these photo monitoring positions throughout the project area. Photos will be taken and compared with the previous photo to assess changes in forest structure and rate of regrowth. Since the longitudinal methods described above require a minimum of 5 years elapsed project time to reveal meaningful changes in forest cover or stocking levels, the project also monitors ongoing activity and event indicators to capture the impact of community mitigation measures. In designing the project strategy community leaders and members identified a number of drivers of deforestation and mitigation measures including: controlling forest fires, closing forests to grazing, closing some forests to fuelwood collection while they regenerate, limiting the conversion of forest lands to quarries and for agriculture, and reducing charcoal making.

Table 20: Ecosystem Service Benefit Indicators

Activity	Activity Indicator (measure annually)	Annual Targets		
		Full Target Achievement	Partial Target Achievement	Missed Target
Fire Control	Number of Hectares Burned during Dry Season by Hima	< 50 ha	50-100	> 100 ha
	Length of fire lines constructed by Hima	> 60 km	40-60 km	< 40 km
Forest Restoration	Number of Hectares with ANR Advance Closure Treatment	> 100 ha	50-100 ha	< 50 ha
	Number of hectares with ANR Silvicultural Treatment	> 50 ha	25-50 ha	< 25 ha
<hr/>				
Impact (after 5 years)	Impact Indicator	Means of assessment	Baseline (2016)	Target (2021)
Forest Condition	Average C-stock in dense forest monitoring plots	Plot measurements	157 tC/ha	200 tC/ha (equivalent to C-stock annual increment of c.8 t/C/ha)
	Average C-stock in open forest monitoring plots	Plot measurements	26 tC/ha	34 tC/ha
Fire damage	Area burnt by wildfires during year	GIS data & project records	64 ha	32 ha

Annually, the activities contributing to REDD+ will be monitored (see Table 20 above). These indicate that the planned REDD+ activities have taken place. Community facilitators (CFs) from each of the 18 micro-watersheds are responsible for collecting this data and reporting the findings to the monitoring officer. The annual monitoring indicator report provides information on changes in carbon stock in the monitoring plots the total area burned by forest fire, and the length of fire lines created to protect forests. This in turn provides an overview of community capacity to limit forest loss and carbon emissions. Annual reporting to the Plan Vivo Foundation will include monitoring results from biomass surveys and photo monitoring for certificate issuance as well as annual activity reports.

10.2. ANR Monitoring

To monitor regeneration in ANR areas, biomass surveys will be carried out annually. At least one plot will be measured and photographed in each ANR area. At least ANR 20 20x20m plots will be established for monitoring purposes over the first three years of the project to assess changes in carbon stock in areas that are being protected by the community through social fencing. In addition, another 20 plots of open forest will be monitored to maintain a baseline. The project also reports on any additional degraded forests that have been placed under “advanced closure” by communities and the area receiving silvicultural forest restoration treatment. Every five years, ANR areas will also be monitored using satellite image analysis as for REDD+. To detect forest regeneration or a lack of change in ANR areas, the perimeters of ANR areas will be marked on maps and satellite images using GPS data

10.3. Environmental & Biodiversity Indicators

The project seeks to address the heavy reliance of project communities on fuelwood by reducing consumption and shifting project families to LPG cooktops. This will take pressure off local forests while improving health conditions within the homes by reduced smoke pollution. Table 21 presents annual indicators to be used to assess project impact. In addition, the project is working with local governments (hima and durbar) to encourage the reduction of area under open pit mining operations. The project will monitor the total area currently being mined in each village to assess how this environmental awareness program is progressing. Finally, the project team will collect data on the observation of key indicator species that are threatened or endangered. Siting data gathered by youth volunteers and community facilitators will be analyzed at the end of each year and included in the annual report to Plan Vivo.

Table 21: Environmental & Biodiversity Indicators

Activity	Activity Indicator (measure annually)	Annual Targets		
		Full Target Achievement	Partial Target Achievement	Missed Target
Fuelwood saving devices	No. of fuel efficient stoves installed	> 50 stoves	25-50 stoves	< 25 stoves
	Number of LPG Units Installed	> 100 units	50-100 units	< 50 units
Biodiversity	Number of biodiversity surveys conducted by CF and youth volunteers	> 2 surveys	1 survey	0 surveys
Quarrying	Number of reports and lobby advocacy meetings reports held	4 reports/ lobbying meetings	2-3 reports/ lobbying meetings	1 or less reports/lobbying meetings
<hr/>				
Impact (after 5 years)	Impact Indicator	Means of assessment	Baseline (2016)	Target (2021)
Fuelwood consumption	Households using fuel efficient stoves (number)	Baseline survey/resurvey	6% of households	At least 25% of all households using fuel efficient stoves
	Households using LPG (number)	Baseline survey/resurvey	1.5% of households	At least 15 % of households using LPG
	Level of household fuelwood consumption (tonnes/year)	Baseline survey/resurvey	2.5 t/yr.	Fuel wood Consumption reduced by an average of 50% across all participating households
Biodiversity	Number of observations of endangered mammal species	Records from surveys conducted by Youth volunteers	42 No. of observation during 2016	50% increase over baseline
Quarrying	% of villages with active quarrying	Baseline assessment	15 % of villages with active quarrying	> 12% of villages with active quarrying

10.4. Socio-Economic Monitoring

The monitoring plan includes socio-economic monitoring to ensure that the project is delivering benefits to participants that enhance their livelihoods and quality of life in accordance with the Plan Vivo Standard. The project seeks to distribute benefits and share them with communities through the provision of annual community development grants (CDG) to each participating village. The village members decide what project they wish to implement and submit proposals to the Federation for funding. Each year, the Federation compiles a report on the type of project, amount spent, and impact of the activity. The Federation also assesses how many community families benefited directly from the project. In addition, the project seeks to build the capacity of community institutions including the Local Working Committees, Self-help groups, and farmer's clubs. Training sessions are held by the Federation to build awareness regarding forest conservation and management, bookkeeping, technical skills in agriculture, animal husbandry, and other income generating activities. The number and results of the trainings are reported each year as an annual indicator (see Table 22 below).

Table 22: Socio-Economic Monitoring Indicators

Activity	Activity Indicator (measure annually)	Annual Targets		
		Full Target Achievement	Partial Target Achievement	Missed Target
Benefit sharing and participation	Number of villages with community Development Grants	> 50 villages	30-50 villages	< 30 villages
	Number of families accessing CDGs	>4000 households	3000-4000 households	< 3000 households
Institutional capacity	Number of training programs	> 10 programs	6-10 programs	< 6 programs
	Number of families participating in income Generating Activities	> 200 families	100-200 families	< 100 families
Impact (after 5 years)	Impact Indicator	Means of assessment	Baseline (2016)	Target (2021)
Knowledge and awareness	Knowledge of the federation & project	Baseline survey/resurvey	75 % of households	85% of all households with knowledge of the Federation and Project activities.
Livelihoods benefits	% of all project households receiving benefits from community grants	Baseline survey/resurvey	30 % of households	60 % of households receiving benefits from community development grants
	% of households with livelihoods activities reflecting conservation of forests and natural resources	Baseline survey/resurvey	20 % of households	60% of all households with expansion of livelihood activities that also reflect conservation of forests and natural resources

10.5. Satellite Monitoring

Satellite image analysis will be carried out as described in Appendix 3. The Federation will use the land use change map to identify areas with reduced rates of deforestation, restoration, and problem areas with continuing deforestation and degradation (see Table 23).

Table 23: Satellite Image Monitoring Indicators

TARGETS	REDD+	ANR
	The rate of deforestation and forest degradation will be assessed by comparing SPOT imagery from the baseline 2010 image with that the most recent images available. Extent of deforestation will be determined by shifts in land use classes, especially between dense, open, and barren areas.	In the ANR areas, rates of forest regrowth will be assessed by comparing forest cover change from the SPOT baseline 2010 period to the most recent image available. Extent of canopy closure (crown cover) will be a primary indicator of forest cover change.
Green target	Reduced by the target percentage or more.	Regeneration in ANR sites is identifiable in the satellite image analysis after year 4.
Amber threshold	Reduced by less than the target percentage, but more than the baseline.	Regeneration is not identifiable in the satellite image analysis after year 4, but a field visit shows some regeneration and there is evidence that an effort has been made to reduce grazing and to implement fire lines.
Red threshold	Same as the baseline or greater.	No change or has been deforested.

10.6. Verification of Targets and Thresholds

REDD+ targets are annual reductions in the rate of deforestation and forest degradation in the project area. In the baseline, dense forest changes to non-forest at a rate of 2.7% per year and dense forest changed to open forest at a rate of 0.1% per year. The annual targets for the reduction in the rate of deforestation and degradation start in 2012 at 33% and increase to 57% by 2021. If there is a reduction in the rate of deforestation and degradation, but it is less than the target, the amber threshold has been met. In this case, the project team and the Community Management Federation will work with communities to improve the implementation of fire lines, sustainable fuel wood collection, reducing charcoal making, reducing grazing, and reducing agricultural expansion.

The ANR target is to achieve identifiable forest regeneration. Forest regeneration should be identifiable in a satellite image analysis after an ANR area has been protected for four years.

If there is regeneration, but there is also evidence of grazing and only partially implemented fire lines, the amber threshold has been met. In this case, the project team and the Community Management Federation will work with communities to improve the implementation of fire lines and to reduce grazing.

11. REFERENCES

Baral et al, 2009. S.K. Baral, R. Malla and S. Ranabhat. Above-ground carbon stock assessment in different forest types of Nepal. *Banko Janakari*, Vol. 19, No. 2.

Brown 1997. Estimating Biomass and Biomass Change of Tropical Forests: a Primer. FAO Forestry Paper – 134. <http://www.fao.org/docrep/w4095e/w4095e00.HTM>

Chave et al, 2005. J. Chave. C. Andalo. S. Brown. M. A. Cairn. J. Q. Chambers. D. Eamus. H. Fořt. Ister. F. Fromard N. Higuchi. T. Kira. J.-P. Lescure. B. W. Nelson H. Ogawa. H. Puig. B. Rie'ra. T. Yamakura. Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Ecosystem Ecology*. *Oecologia* (2005) 145: 87-99.

FSI 1996. Volume Equations for Forests of India, Nepal and Bhutan”, Published by: Forest Survey of India, Ministry of Environment & Forests, Govt. of India, (1996)

FSI 2006. Forest Survey of India. State of Forest Report. FSI, Dehradun.

Jina et al, 2008. B.S. Jina, Pankaj Sah, M.D. Bhatt and Y.S. Rawat. Estimating carbon sequestration rates and total carbon stockpile in degraded and non-degraded sites of oak and pine forest of Kumaun Central Himalaya. *Ecoprint* 15: 75-81, 2008 ISSN 1024-8668. Ecological Society (ECOS), Nepal.

Meghalaya Tourism, 2012. Department of Tourism, Government of Meghalaya <http://megtourism.gov.in/ecodestination.html>. Accessed April 2012.

Shrestha, R. 2010. Participatory carbon monitoring: An experience from the Koshi Hills, Nepal.

Ravi K Shrestha, Programme Officer, Livelihoods and Forestry Programme, Dhankuta. Nepal.
ravikshrestha@gmail.com

Rawat, L., Kamboj, S.K., and Kandwal, A. 2015. Biomass Expansion Factor and Root-to-Shoot Ratio of Some Tree Species of Punjab, India. *Indian Forester*, [S.I.], p. 146-153, feb. 2015. ISSN 2321-094X. Available at:
<<http://www.indianforester.co.in/index.php/indianforester/article/view/60650>>

Zanne 2009. Zanne, A.E., Lopez-Gonzalez, G., Coomes, D.A., Ilic, J., Jansen, S., Lewis, S.L., Miller, R.B., Swenson, N.G., Wiemann, M.C., and Chave, J. (2009) Global wood density database. Dryad. <http://hdl.handle.net/10255/dryad.235>

APPENDICES

Appendix 1: Mawphlang Pilot Project

Pilot project activities have been successful in Mawphlang. From 2006 to 2010, REDD+ and ANR activities have taken place in Mawphlang. Fire has been reduced by fire lines and fire watches, and assisted natural regeneration has led to forest regrowth.

