

**SAFETY**

Wear suitable gloves, overalls and face shield when handling this product.

**STORAGE**

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Slight thickening can be a product characteristic under extreme conditions.

**DISPOSAL**

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

**WARRANTY**

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

**COMPATABILITY**

**Mn – Cu SC** is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.

**INSTRUCTIONS FOR USE**

Always shake the bottle well before use.

**NEVER MIX PESTICIDES IN THEIR CONCENTRATED FORM.**

If Spray volumes are less than 200l/ha or if you are in doubt, then perform a "Jar test" on the mix before starting.

Half fill the tank with water and switch on agitation. Always make sure the additions are fully mixed until dissolved or dispersed before adding the next product.

- Add water conditioners, if needed. Add granules powders and mix well. Pre disperse if possible (WP, DG).
- Add **Mn – Cu SC** and other flowables (SC, SE).
- Add Emulsion Concentrates and crop oils (EC).
- Add water soluble: liquids and solids. (SL, SP).
- Add wetters, if needed and top up with water.

Provided these guidelines are followed almost all tank mixes are physically compatible, with continuous agitation. If agitation fails, for any reason, then it is essential to clean out the tank immediately.

**APPLICATION RECOMMENDATIONS**

CROP	RATE	TIMING
Apply the product in a minimum of 200 litre of water unless specified		
Carrots	1l/ha	When crop is 15cm tall. In case of need, repeat at 10–14 day intervals.
Cereals	1l/ha	From 2 leaves to second node detectable (Zadok's GS 12 to 32) Where required repeat at 7–14 day interval.
Maize	1l/ha	Apply at 4–8 leaves.
Peas, Beans	1l/ha	At 4–6 leaves 'Marsh Spot'.
Potatoes	1l/ha	One week after 100% emergence. Repeat if necessary at intervals of 10–14 days.
Sugar Beet	1l/ha	From 4–6 leaves.
Additional Crops: For additional crops not listed here, check our website at <a href="http://WWW.SAFAGROW.COM">WWW.SAFAGROW.COM</a>		



**WARNING**

EUH208:	Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction.
H319:	Causes serious eye irritation.
H411:	Toxic to aquatic life with long lasting effects.
P264:	Wash hands thoroughly after handling.
P273:	Avoid release to the environment.
P280:	Wear protective gloves/protective clothing/eye protection/face protection.
P305+351+338:	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P337+313:	If eye irritation persists: Get medical attention.
P391:	Collect spillage.
UFI: VQ31-YQKW-R00N-2TU2	

VAL1065BUK

**EC FERTILISER**

Micro-nutrient fertiliser	
Copper (Cu) Total	5.7%w/w
5.7%w/w Oxchloride	
Manganese (Mn) Total	22.7%w/w
22.7%w/w Carbonate	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	