

SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Crystallisation can be a product characteristic under extreme conditions.

DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

INSTRUCTIONS FOR USE

Read the label before use.

Shake the bottle well before use.

Half fill spray tank with water and begin agitation. Empty the required quantity into the spray tank, mix until dissolved or dispersed and add the balance of the water. Ensure agitation is maintained throughout the spraying operation. Do not allow the spray tank agitation to stop whilst product remains in the tank.

COMPATABILITY

Bio + is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.

APPLICATION RECOMMENDATIONS

CROP	RATE	TIMING
Apple	2.5 ml/l	At pea stage
Avocado	2.5 ml/l	Apply from flowering and then every 14 days
Beans	2.0 ml/l	Apply from flowering and then every 14 days
Brussel Sprouts	2.5 ml/l	Apply from flowering and then every 14 days
Banana	2.0 ml/l	Apply every 20 days from formation of fruit
Cabbage	2.0 ml/l	At button size head formation
Cherry, Plums	2.0 ml/l	Apply at the formation of pea sized fruit and then every 14 days
Cotton	2.0 ml/l	Apply at 4 leaf stage and at flower initiation
Peppers, Tomatoes	2.0 ml/l	Apply from flowering and then every 14 days
Potatoes	2.5 ml/l	Apply between 40–45 days after planting
Strawberries	2.0 ml/l	Apply from the 40th day after planting followed by three applications every 2 weeks
Pears	2.5 ml/l	Apply from petal fall with 3–4 applications every 14 days
Paddy	2.5 ml/l	Apply at panicle formation
Carnation	2.0 ml/l	Apply every 15 days from 7 days after first pinching
Gerbera	2.0 ml/l	Apply twice a month from 60 days after first pinching
Rose	2.0 ml/l	Apply twice a month to both old and new plants
Nursery	2.0 ml/l	Apply at 4 leaf stage and repeat after 7 days
Additional Crops: For additional crops not listed here, check our website at WWW.SAFAGROW.COM		

EC FERTILISER

NPK – fertiliser suspension 12.6 – 12.6 – 12.6 with micro-nutrients	
Nitrogen (N) Total	12.6%w/w
4.5%w/w Nitric nitrogen	
3.9%w/w Ammoniacal nitrogen	
4.2%w/w Ureic nitrogen	
Phosphorus Pentoxide (P ₂ O ₅) Water-soluble	12.6%w/w
Potassium Oxide (K ₂ O) Water-soluble	12.6%w/w
Boron (B) Water-soluble	0.020%w/w
Copper (Cu) Water-soluble	0.047%w/w
0.047%w/w chelated by EDTA	
Iron (Fe) Water-soluble	0.152%w/w
0.152%w/w chelated by EDTA	
Manganese (Mn) Water-soluble	0.047%w/w
0.047%w/w chelated by EDTA	
Molybdenum (Mo) Water-soluble	0.001%w/w
Zinc (Zn) Water-soluble	0.047%w/w
0.047%w/w chelated by EDTA	
Chelated fraction stable within pH range 3–10	

H412:	Harmful to aquatic life with long lasting effects.
P273:	Avoid release to the environment.
P501:	Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.
UFI: 3TC6-N015-P00X-WF0N	

VAL1215BUK