

#### SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

#### STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Crystallisation can be a product characteristic under extreme conditions.

#### DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

#### WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

#### INSTRUCTIONS FOR USE

Read the label before use.

Shake the bottle well before use.

Half fill spray tank with water and begin agitation. Empty the required quantity into the spray tank, mix until dissolved or dispersed and add the balance of the water. Ensure agitation is maintained throughout the spraying operation. Do not allow the spray tank agitation to stop whilst product remains in the tank.

#### COMPATABILITY

**Max K** is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.



H411:	Toxic to aquatic life with long lasting effects.
P273:	Avoid release to the environment.
P391:	Collect spillage.
P501:	Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.
UFI: F5G7-303Y-600W-H03F	

VAL1355BUK

#### APPLICATION RECOMMENDATIONS

CROP	RATE	TIMING
Cereals	2.0 ml/l	For potassium deficient crops apply every 14 days
Vegetables	2.5 ml/l	Apply from fruiting and then every 14 days
Cabbage, Cauliflower	2.0 ml/l	Apply at button size head formation and then every 14 days
Cherry, Plums	2.0 ml/l	Apply at the formation of pea sized fruit and then at 14 day intervals
Peppers, Tomatoes, Brinjals, Cucurbits	2.0 ml/l	Apply at flowering and then every 14 days
Potatoes	2.5 ml/l	Apply between 40–45 days after planting
Ornamentals, Greenhouse, Polyhouse	2.0 ml/l	Apply twice a month from 45–50 days after planting
Additional Crops: For additional crops not listed here, check our website at <a href="http://WWW.SAFAGROW.COM">WWW.SAFAGROW.COM</a>		

#### EC FERTILISER

NPK – fertiliser suspension 6.1 – 12.2 – 24.4 with micro-nutrients	
Nitrogen (N) Total	6.1%w/w
0.6%w/w Nitric nitrogen	
0.8%w/w Ammoniacal nitrogen	
4.7%w/w Ureic nitrogen	
Phosphorus Pentoxide (P <sub>2</sub> O <sub>5</sub> ) Water-soluble	12.2%w/w
Potassium Oxide (K <sub>2</sub> O) Water-soluble	24.4%w/w
Boron (B) Water-soluble	0.021%w/w
Copper (Cu) Water-soluble	0.054%w/w
Iron (Fe) Water-soluble	0.107%w/w
0.107%w/w chelated by EDTA	
Manganese (Mn) Water-soluble	0.054%w/w
Molybdenum (Mo) Water-soluble	0.001%w/w
Zinc (Zn) Water-soluble	0.054%w/w
Chelated fraction stable within pH range 3–10	