



Mn – Zn SC



Wear suitable gloves, overalls and face shield when handling this product.

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Slight thickening can be a product characteristic under extreme conditions.

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

INSTRUCTIONS FOR USE

Read the label before use

Shake the bottle well before use.

Half fill spray tank with water and begin agitation. Empty the required quantity into the spray tank, mix until dissolved or dispersed and add the balance of the water. Ensure agitation is maintained throughout the spraying operation. Do not allow the spray tank agitation to stop whilst product remains in the tank.

COMPATABILITY

Mn - Zn SC is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.





DANGER

EUH208:	Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction.		
H314:	Causes severe skin burns and eye damage.		
H400:	Very toxic to aquatic life.		
H411:	Toxic to aquatic life with long lasting effects.		
P260:	Do not breathe mist.		
P264:	Wash hands thoroughly after handling.		
P273:	Avoid release to the environment.		
P280:	Wear protective gloves.		
P305+351+338:	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.		
P391:	Collect spillage.		

VAL1315BUK

APPLICATION RECOMMENDATIONS

CROP				
Apply the product in a minimum of 200 litre of water unless specified				
Carrots	1l/ha	When crop is 15cm tall. In case of need, repeat at 10–14 day intervals.		
Cereals	1I/ha	From 2 leaves to second node detectable (Zadok's GS 12 to 32) Where required repeat at 7–14 day interval.		
Maize	1l/ha	Apply at 4–8 leaves.		
Peas, Beans	1l/ha	At 4–6 leaves 'Marsh Spot'.		
Potatoes	1l/ha	One week after 100% emergence. Repeat if necessary at intervals of 10–14 days.		
Sugar Beet	1l/ha	From 4–6 leaves.		
Additional Committee of the second se				

Additional Crops: For additional crops not listed here, check our website at WWW.SAFAGROW.COM

14.4%w/w
19.0%w/w

To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

