



**SAFETY**

Wear suitable gloves, overalls and face shield when handling this product.

**STORAGE**

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Slight thickening can be a product characteristic under extreme conditions.

**DISPOSAL**

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

**WARRANTY**

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

**COMPATIBILITY**

Zinc Alloy® is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.

**INSTRUCTIONS FOR USE**

Always shake the bottle well before use.

NEVER MIX PESTICIDES IN THEIR CONCENTRATED FORM.

If Spray volumes are less than 200l/ha or if you are in doubt, then perform a "Jar test" on the mix before starting.

Half fill the tank with water and switch on agitation. Always make sure the additions are fully mixed until dissolved or dispersed before adding the next product.

- Add water conditioners, if needed. Add granules powders and mix well. Pre disperse if possible (WP, DG).
- Add Zinc Alloy® and other flowables (SC, SE).
- Add Emulsion Concentrates and crop oils (EC).
- Add water soluble: liquids and solids. (SL, SP).
- Add wetters, if needed and top up with water.

Provided these guidelines are followed almost all tank mixes are physically compatible, with continuous agitation. If agitation fails, for any reason, then it is essential to clean out the tank immediately.

**APPLICATION RECOMMENDATIONS**

CROP	RATE	TIMING
Apply the product in a minimum of 200 litre of water unless specified		
Brassica	1l/ha	At 4-9 leaves.
Carrots	1l/ha	When crop is 15 cm tall. In case of moderate to severe deficiency repeat at 10 to 14 day intervals.
Cereals	1l/ha	From 2 leaves to first node detectable (Zadok's G.S. 12-31).
Hops	1.5l/ha	When 2.5M tall Water rate: 500 l/ha
Maize	1l/ha	From 3 to 8 leaves. In case of severe deficiency repeat applications in intervals of 10-14 days.
Potatoes	1l/ha	One week after 100% emergence and repeated as necessary in intervals of 10-14 days in case of moderate to severe deficiency.
Sugar beet	1l/ha	From 4-6 leaf stage. Repeat applications in intervals of 10-14 days where required.
Additional Crops: For additional crops not listed here, check our website at <a href="http://WWW.SAFAGROW.COM">WWW.SAFAGROW.COM</a>		



**WARNING**

H315:	Causes skin irritation.
H319:	Causes serious eye irritation.
H410:	Very toxic to aquatic life with long lasting effects.
P264:	Wash hands, forearms and face thoroughly after handling.
P273:	Avoid release to the environment.
P280:	Wear protective clothing, eye protection, face protection.
P302+P352:	IF ON SKIN: Wash with plenty of water.
P305+P351+P338:	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P391:	Collect spillage.
UFI: MHFD-404Q-W00P-VSUF	

**EC FERTILISER**

Zinc fertiliser in suspension	
Zinc (Zn) Total	38.7%/w
37.1%/w Oxide	
1.6%/w Nitrate	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	