



### SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

### STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Slight thickening can be a product characteristic under extreme conditions.

### DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

### WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

### INSTRUCTIONS FOR USE

Read the label before use.

Shake the bottle well before use.

Half fill spray tank with water and begin agitation. Empty the required quantity into the spray tank, mix until dissolved or dispersed and add the balance of the water. Ensure agitation is maintained throughout the spraying operation. Do not allow the spray tank agitation to stop whilst product remains in the tank.

### COMPATABILITY

**Calcium Aloy** is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.

### APPLICATION RECOMMENDATIONS

CROP	RATE	TIMING
Apply the product in a minimum of 200 litre of water unless specified		
Apples and Pears	3–5l/ha	Apply and repeat at 10 day intervals from 4 weeks after petal fall. Water rate 500 L/Ha minimum.
Broccoli, Cabbage, Calabrese, cauliflower and sprouts	3–5l/ha	Apply from 4–9 leaves and repeat at 10–14 day intervals.
Carrots	3–5l/ha	In case of moderate to severe deficiency repeat same rate at 10 to 14 day intervals.
Cucurbits (field grown)	3–5l/ha	Apply and repeat 1 7 day intervals during fruit development.
Lettuce (field grown)	3–5l/ha	Apply 10–14 days from transplanting or emergence and repeat as required in intervals of 10–14 days.
Onions (field grown)	3–5l/ha	Apply 10–14 days from transplanting or emergence and repeat as required in intervals of 10–14 days.
Potatoes	5l/ha	Apply 14 days after emergence and repeat as necessary at the same rate in intervals of 10–14 days.
Additional Crops: For additional crops not listed here, check our website at <a href="http://WWW.SAFAGROW.COM">WWW.SAFAGROW.COM</a>		



## WARNING

H319:	Causes serious eye irritation.
H412:	Harmful to aquatic life with long lasting effects.
P264:	Wash hands thoroughly after handling.
P273:	Avoid release to the environment.
P280:	Wear protective gloves/protective clothing/eye protection/face protection.
P305+351+338:	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P337+313:	If eye irritation persists: Get medical attention.
P501:	Dispose of contents/container to hazardous or special waste collection point.

VAL1565BUK

### EC FERTILISER

Micro-nutrient fertiliser	
Manganese (Mn) Total	1.5%w/w
1.5%w/w Nitrate	
Zinc (Zn) Total	0.6%w/w
0.6%w/w Oxide	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	