

SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Slight thickening can be a product characteristic under extreme conditions.

DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

COMPATABILITY

Mn – Cu – Zn SC is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.

INSTRUCTIONS FOR USE

Always shake the bottle well before use.

NEVER MIX PESTICIDES IN THEIR CONCENTRATED FORM.

If Spray volumes are less than 200l/ha or if you are in doubt, then perform a "Jar test" on the mix before starting.

Half fill the tank with water and switch on agitation. Always make sure the additions are fully mixed until dissolved or dispersed before adding the next product.

- Add water conditioners, if needed. Add granules powders and mix well. Pre disperse if possible (WP, DG).
- Add **Mn – Cu – Zn SC** and other flowables (SC, SE).
- Add Emulsion Concentrates and crop oils (EC).
- Add water soluble: liquids and solids. (SL, SP).
- Add wetters, if needed and top up with water.

Provided these guidelines are followed almost all tank mixes are physically compatible, with continuous agitation. If agitation fails, for any reason, then it is essential to clean out the tank immediately.

APPLICATION RECOMMENDATIONS

CROP	RATE	TIMING
Apply the product in a minimum of 200 litre of water unless specified		
Carrots	1l/ha	When crop is 15cm tall. In case of need, repeat at 10–14 day intervals.
Cereals	1l/ha	From 2 leaves to second node detectable (Zadok's GS 12 to 32) Where required repeat at 7–14 day interval.
Maize	1l/ha	14 days after transplanting or where direct sown, when the crop is 15 cm tall.
Peas, Beans	1l/ha	At 4–6 leaves 'Marsh Spot'.
Potatoes	1l/ha	One week after 100% emergence. Repeat if necessary at intervals of 10–14 days.
Sugar Beet	1l/ha	From 4–6 leaves.
Additional Crops: For additional crops not listed here, check our website at WWW.SAFAGROW.COM		



WARNING

EUH208:	Contains 1,2-benzisothiazolin-3-one(2634-33-5). May produce an allergic reaction.
H302:	Harmful if swallowed.
H319:	Causes serious eye irritation.
H410:	Very toxic to aquatic life with long lasting effects.
P264:	Wash hands, forearms and face thoroughly after handling.
P270:	Do not eat, drink or smoke when using this product.
P280:	Wear protective clothing, eye protection, face protection.
P301+P312:	IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell.
P305+P351+P338:	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P391:	Collect spillage.
UFI: CCX1-90YM-A00M-VW2C	

VAL1105BUK

EC FERTILISER

Micro-nutrient fertiliser	
Copper (Cu) Total	6.3%w/w
6.3%w/w Oxychloride	
Manganese (Mn) Total	18.8%w/w
18.8%w/w Carbonate	
Zinc (Zn) Total	4.8%w/w
4.8%w/w Oxide	
To be used only where there is a recognised need. Do not exceed the appropriate dose rates.	