

SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Crystallisation can be a product characteristic under extreme conditions.

DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

INSTRUCTIONS FOR USE

Read the label before use.

Shake the bottle well before use.

Half fill spray tank with water and begin agitation. Empty the required quantity into the spray tank, mix until dissolved or dispersed and add the balance of the water. Ensure agitation is maintained throughout the spraying operation. Do not allow the spray tank agitation to stop whilst product remains in the tank.

COMPATABILITY

MAX Ca is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.



WARNING

H302:	Harmful if swallowed.
P264:	Wash hands, forearms and face thoroughly after handling.
P270:	Do not eat, drink or smoke when using this product.
P301+P312:	IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell.
P330:	Rinse mouth.
P501:	Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.
UFI: HR3Y-E1AE-F009-7SKT	

VAL1225BK

APPLICATION RECOMMENDATIONS

CROP	RATE	TIMING
Apples	2.5 ml/l	Apply from Petal fall with 3–4 applications every 14 days
Avocado	2.5 ml/l	From flowering and then every 14 days
Beans	2.0 ml/l	From flowering and then every 14 days
Brussel Sprouts	2.5 ml/l	From flowering and then every 14 days
Banana	2.0 ml/l	Apply every 20 days from fruit formation
Cherry, Plums	2.0 ml/l	Apply at formation of pea sized fruit and then every 14 days
Cotton	2.0 ml/l	Apply at flower initiation
Peppers, Tomatoes, Brinjals, Cucurbits	2.0 ml/l	Apply at flowering and then every 14 days
Potatoes	2.5 ml/l	Apply 40–45 days after planting
Strawberries	2.0 ml/l	Apply from the 40th day after planting followed by 3 further applications every 14 days
Pears	2.5 ml/l	Apply at petal fall followed by a further 3 to 4 applications every 14 days
Paddy	2.5 ml/l	Apply at panicle formation
Carnation	2.0 ml/l	Apply every 15 days from 7 days after first pinching
Gerbera	2.0 ml/l	Apply twice a month from 60 days after planting
Rose	2.0 ml/l	Apply twice a month to both old and new plants
Additional Crops: For additional crops not listed here, check our website at WWW.SAFAGROW.COM		

EC FERTILISER

Nitrogen (N) Total	9.5% w/w
8.4% w/w Nitric nitrogen	
1.1% w/w Ureic nitrogen	
Calcium Oxide (CaO) Water-soluble	14.2% w/w
Boron (B) Water-soluble	0.047% w/w
Copper (Cu) Water-soluble	0.019% w/w
0.019% w/w chelated by EDTA	
Iron (Fe) Water-soluble	0.047% w/w
0.047% w/w chelated by EDTA	
Manganese (Mn) Water-soluble	0.095% w/w
0.095% w/w chelated by EDTA	
Molybdenum (Mo) Water-soluble	0.001% w/w
Zinc (Zn) Water-soluble	0.019% w/w
0.019% w/w chelated by EDTA	
Chelated fraction stable within pH range 1–7	