

1. A particle P moves on the x -axis. The acceleration of P at time t seconds, $t \geq 0$, is $(3t + 5) \text{ m s}^{-2}$ in the positive x -direction. When $t = 0$, the velocity of P is 2 m s^{-1} in the positive x -direction. When $t = T$, the velocity of P is 6 m s^{-1} in the positive x -direction. Find the value of T .

(6)




(Total 6 marks)





(Total 9 marks)



(4)

(b) Find the value of a .

(4)

Question 4 continued




(Total 8 marks)







(Total 9 marks)






Figure 2 shows a uniform rod AB of mass m and length $4a$. The end A of the rod is freely hinged to a point on a vertical wall. A particle of mass m is attached to the rod at B . One end of a light inextensible string is attached to the rod at C , where $AC = 3a$. The other end of the string is attached to the wall at D , where $AD = 2a$ and D is vertically above A . The rod rests horizontally in equilibrium in a vertical plane perpendicular to the wall and the tension in the string is T .

- (5)

(3)






Question 6 continued



(Total 8 marks)





A ball is projected with speed 40 ms^{-1} from a point P on a cliff above horizontal ground. The point O on the ground is vertically below P and OP is 36 m. The ball is projected at an angle θ° to the horizontal. The point Q is the highest point of the path of the ball and is 12 m above the level of P . The ball moves freely under gravity and hits the ground at the point R , as shown in Figure 3. Find

- (a) the value of θ , (3)
- (b) the distance OR , (6)
- (c) the speed of the ball as it hits the ground at R . (3)





Q7







